

Preventing HIV Transmission in Persons with HIV

This is a PDF version of the following document:

Module 5: [Prevention of HIV](#)

Lesson 2: [Preventing HIV Transmission in Persons with HIV](#)

You can always find the most up-to-date version of this document at

<https://www.hiv.uw.edu/go/prevention/prevention-positives/core-concept/all>.

Background

The availability of widespread effective antiretroviral therapy has transformed HIV from a fatal infection to a manageable chronic disease. Despite major advances in antiretroviral therapy, the incidence of new HIV infections in the United States continues to occur with more than 30,000 annual new HIV infections ([Figure 1](#)).^[1] For persons living with HIV, antiretroviral therapy can dramatically reduce HIV transmission to others.^[2,3,4,5] All health professionals involved with HIV care should have knowledge and awareness of effective strategies that can play a role in reducing HIV transmission. The use of antiretroviral therapy in people with HIV has emerged as a highly impactful means to prevent HIV transmission and this approach is referred to as Treatment as Prevention (TasP).^[6] This topic review will focus on the biomedical, behavioral, and structural measures related to preventing transmission from persons with HIV, with an emphasis on Treatment as Prevention. The discussion of preventing HIV acquisition among individuals who do not have HIV is discussed in detail in the lesson [HIV Preexposure Prophylaxis](#).

Antiretroviral Treatment as Prevention

HIV RNA Level and Risk of HIV Transmission

A sentinel study in Rakai, Uganda, first reported HIV RNA levels tightly correlated with the risk of heterosexual HIV transmission.^[7] This study involved 415 serodifferent heterosexual couples not taking antiretroviral therapy.^[7] The main finding was that mean serum HIV RNA was significantly higher in subjects whose partners seroconverted than in subjects whose partners remained HIV negative, and HIV transmission was considered rare among persons who had an HIV RNA level less than 1,500 copies/mL (Figure 2).^[7] This study played a major role in spurring on studies that would examine the impact that lowering HIV RNA levels with antiretroviral therapy would have on sexual transmission of HIV.

Antiretroviral Therapy in HIV Serodifferent Couples

Convincing data from several studies have shown that antiretroviral treatment taken by individuals with HIV dramatically reduces HIV transmission to their sex partners.^[3,4,8,9]

- **HPTN-052:** The concept of treating persons with HIV to prevent transmission to others, often referred to as treatment as prevention, garnered major support following the release of data from the landmark HPTN-052 trial.^[3,4] The HPTN-052 trial was a randomized, controlled study that enrolled 1,763 HIV serodifferent, predominantly heterosexual couples from 9 countries. All persons with HIV had a CD4 count of 350 to 550 cells/mm³ at enrollment, and none had HIV-related symptoms. The trial demonstrated that early initiation of antiretroviral therapy (started at the time of enrollment) reduced rates of sexual transmission of HIV to the partner without HIV by 96%, when compared with deferral of antiretroviral therapy (started when the CD4 count decreased to less than 250 cells/mm³ or at the onset of an AIDS-related event) (Figure 3).^[3,4]
- **PARTNER-1 Study:** In the first phase of the European PARTNER (Partners of People on ART—A New Evaluation of the Risks) study, investigators at 75 sites in 14 European countries evaluated the impact of antiretroviral therapy on HIV transmission risk in 888 HIV-serodifferent couples engaging in condomless sex, including 548 heterosexual couples and 340 gay male couples.^[2] The eligibility for enrollment required the partner with HIV to be taking antiretroviral therapy and have an HIV RNA level less than 200 copies/mL.^[2] Enrollment took place from September 2010 through May 2014, and during this time, there were zero phylogenetically-linked HIV transmissions that occurred in these couples, with an estimated 58,000 condomless sex acts (22,000 in gay male couples and 36,000 in heterosexual couples).^[2] There were 11 new HIV infections during the study period, but none of these were phylogenetically linked.
- **PARTNER-2 Study:** In the second phase of the European PARTNER (Partners of People on ART—A New Evaluation of the Risks) study, investigators at the same 75 sites in 14 European countries continued the PARTNER study, but additional enrollment was only for serodifferent gay men.^[10] The enrollment period analyzed for PARTNER-2 was September 2010 through July 31, 2017, and included a total of 972 HIV serodifferent gay male couples.^[10] As with PARTNER-1, the eligibility for enrollment required the partner with HIV to be taking antiretroviral therapy and have an HIV RNA level less than 200 copies/mL. During the study, the serodifferent gay male couples reported condomless anal sex a total of 76,088 times, and there were zero phylogenetically-linked transmissions.^[10] There were, however, 15 new HIV infections during the study period, but none of these were phylogenetically linked.^[10]
- **Opposites Attract:** The Opposites Attract trial was conducted from 2012 through 2016 and enrolled 358 HIV-serodifferent gay male couples in Thailand, Brazil, and Australia.^[11] Of the 358 couples that enrolled, 343 had at least one follow-up visit, and 75% (258 of 343) of the partners with HIV had an HIV RNA level less than 200 copies/mL.^[11] There were zero phylogenetically-linked cases of HIV transmission from among 12,447 sex acts that involved (1) condomless anal intercourse, (2) partners with HIV taking antiretroviral therapy (and with an HIV RNA less than 200 copies/mL), and (3) partners without HIV not taking HIV preexposure prophylaxis (PrEP).^[11] There were three new HIV infections

during the study period, but none were phylogenetically linked.[11]

Undetectable Equals Untransmittable

Extensive data from multiple studies strongly support the concept that persons with HIV who consistently take antiretroviral therapy and maintain undetectable HIV RNA levels do not transmit HIV sexually to others, even with condomless sex.[2,10,11] This concept is commonly referred to as undetectable equals untransmittable.[5] The undetectable equals untransmittable concept is an extremely important message that is now widely endorsed by prominent scientists, clinicians, organizations, and societies.[5] The findings from the studies listed above serve as the foundation for the undetectable equals untransmittable concept and these studies underscore the tremendous impact that antiretroviral treatment can have in preventing transmission of HIV from persons with HIV.[2,10,11] These findings also emphasize the benefit of routine HIV testing and prompt initiation of antiretroviral therapy for persons who acquire HIV. For persons with HIV who are having condomless sex, it is important to perform regular screening for sexually transmitted infections, since antiretroviral therapy does not provide protection against common sexually transmitted infections, such as chlamydia, gonorrhea, and syphilis.

Recommendations for Antiretroviral Treatment as Prevention

The Adult and Adolescent ARV Guidelines cite prevention of sexual transmission of HIV as one of the main reasons to recommend antiretroviral therapy for all persons with HIV (Treatment as Prevention).[6,12] Further, all people with HIV should be informed that maintaining a plasma HIV RNA level less than 200 copies/mL with antiretroviral therapy will prevent sexual transmission of HIV.[12] Note, these recommendations do not address use of antiretroviral treatment to prevent HIV transmission among persons who inject drugs, but theoretically, use of antiretroviral therapy with achievement of undetectable plasma HIV RNA levels should decrease bloodborne transmission of HIV. The table below summarizes recommendations regarding the use of antiretroviral therapy to prevent sexual transmission of HIV ([Table 1](#)).[6]

Knowledge of HIV Status

The proportion of persons who are aware of their HIV has steadily increased from an estimated 75% in 2003 to 87% in 2022.[[1](#),[13](#),[14](#)] Several HIV transmission modeling studies for the United States have concluded that persons unaware of their HIV diagnosis account for a disproportionately higher number of transmitted HIV infections than persons aware of their HIV diagnosis.[[15](#),[16](#),[17](#)] In the 2016 CDC Progression and Transmission of HIV (PATH 2.0) model, investigators estimated that among all persons with HIV in the United States in 2016, the 14.5% of persons with undiagnosed HIV accounted for 37.5% of new HIV transmissions that year ([Figure 4](#)).[[17](#)] Knowledge of HIV serostatus is the first step toward linking persons to HIV medical care whereby they can receive antiretroviral therapy. Individuals with HIV who are unaware of their status are not taking antiretroviral therapy and thus are missing out on this critical element of HIV prevention. In the Ending HIV Epidemic—A Plan for the United States, diagnosing all individuals with HIV as early as possible after infection is one of the main pillars in the strategic initiative.[[18](#)]

Behavioral Prevention Interventions

Risk Reduction Counseling

Counseling to reduce activities that can increase the risk of HIV transmission to others is inadequate as a primary strategy for reducing HIV transmission. For persons with HIV, the impact of consistently taking antiretroviral therapy and maintaining undetectable HIV RNA levels far exceeds the impact of prevention strategies that rely on behavioral interventions. Nevertheless, risk reduction counseling for persons with known HIV remains a complementary piece of a comprehensive prevention strategy.^[19] The CDC has identified evidence-based risk reduction counseling strategies for people with HIV.^[20] Moreover, older studies that examined the impact of behavioral interventions for people with HIV showed a reduction in self-reported condomless sex as well as a decline in the incidence of sexually transmitted infections in persons who received behavioral interventions.^[21]

HIV Status Disclosure and Partner Testing

Partner counseling and referral services is a public health service that helps people with HIV disclose their HIV status to current or former sex or injection drug partners. The public health system provides a trained counselor who can work with the person newly diagnosed with HIV to support disclosure to partners, as well as to provide partner notification directly in cases where the person with newly diagnosed HIV is not able to disclose their HIV status. Partner notification and testing is important because of the high yield in HIV case finding.^[22,23] In two national studies, partner counseling and referral services, including partner notification and HIV testing, effectively identified a substantial number of partners with a new HIV diagnosis; in these studies, 8% of the partners of persons newly diagnosed with HIV tested positive for HIV, and these results were consistent across a 10-year period.^[24,25]

Condom Use

Extensive data has shown that persons with HIV who are taking antiretroviral therapy and maintain HIV RNA levels consistently less than 200 copies/mL do not transmit HIV sexually to others, even with condomless sex.^[2,5,10,11] Accordingly, efforts emphasizing condom use as the primary method for HIV prevention have markedly diminished in recent years. Nevertheless, condoms still have a role for persons with HIV, especially in persons newly starting on antiretroviral therapy and for persons who do not have consistently suppressed HIV RNA levels. In addition, condoms play a key role in preventing the transmission and acquisition of other sexually transmitted infections. Consistent, correct condom use decreases HIV transmission by 70 to 80% among HIV-serodifferent heterosexual couples when compared with non-condom users.^[26,27,28] In a separate analysis of the protective effect of condom use among HIV-serodifferent male couples who have anal sex, consistent condom use reduced the risk of HIV transmission by approximately 70%.^[29]

Serosorting and Seropositioning

The practice of serosorting and seropositioning are self-selected behaviors intended to reduce HIV transmission risk and are referred to as seroadaptive strategies.^[30,31] Serosorting describes the practice of choosing sex partners based on concordant HIV status, typically with the practice of selectively using condoms only when sex occurs with persons of a serodifferent HIV status. Data on the impact of serosorting have been mixed.^[32,33,34] Seropositioning refers to choosing a different sexual position or practice based on the HIV serostatus of one's partner. For men utilizing seropositioning, the person with HIV typically takes the receptive role during unprotected anal sex when the partner with HIV has the insertive role. There are no guidelines in the United States that recommend serosorting or seropositioning as an impactful prevention measure. In addition, the interest in serosorting and seropositioning as an HIV prevention measure has markedly decreased with the emphasis on HIV treatment as prevention and HIV preexposure prophylaxis (PrEP).

Importance of Diagnosing and Treating Acute HIV

At the time of early HIV infection (less than 6 months after HIV acquisition), patients usually have high HIV RNA levels and lack significant neutralizing antibodies and thus are considered highly infectious.[[35,36,37](#)] Studies have evaluated the relative likelihood of transmitting HIV during acute or early HIV infection, and estimate a significantly higher risk of transmission with acute (or early) HIV than with chronic HIV infection.[[38,39,40,41](#)] Indeed, using the 2016 CDC Progression and Transmission of HIV (PATH 2.0) model, the Centers for Disease Control and Prevention estimated the highest risk of HIV transmission along the HIV continuum of care occurred in persons with acute HIV who were unaware of their HIV diagnosis ([Figure 6](#)).[[17](#)] Accordingly, it is important to diagnose individuals with acute and recent (early) HIV whenever possible. To this end, the CDC HIV testing algorithm recommends the use of the HIV-1/2 antigen-antibody immunoassay for use as the initial HIV screening test in an effort to improve the diagnosis of persons with acute HIV.[[42,43](#)] In addition, all persons diagnosed with acute HIV should immediately start antiretroviral therapy to reduce the risk of HIV transmission, as well as to garner potential long-term immunologic benefit from early therapy.[[44,45](#)]

Screening and Treating Sexually Transmitted Infections

Sexually transmitted infections can facilitate transmission and acquisition of new HIV infection, especially when the individual with HIV is not taking suppressive antiretroviral therapy.[46] Rates of sexually transmitted infections, particularly syphilis, gonorrhea, and chlamydia are disproportionately higher among men with HIV who have sex with men.[47] Non-injection drug use, particularly with methamphetamines, as well as recreational use of erectile-enhancing medications, among men with HIV who have sex with men has been implicated in transmission of sexually transmitted infections.[48,49] Available data on herpes simplex virus (HSV) has not shown a convincing reduction in HIV transmission risk with acyclovir suppressive therapy taken by persons with HIV.[50] Screening, diagnosis, and treatment of sexually transmitted infections in persons with HIV remain a priority, but in the current era, the impact of treatment of sexually transmitted infections on preventing HIV transmission is unclear, especially for persons taking fully suppressive antiretroviral therapy.

- **Screening Recommendations for Sexually Transmitted Infections:** The 2021 STI Treatment Guidelines recommend that all sexually active persons with HIV undergo routine screening for sexually transmitted infections at all exposed anatomic sites (e.g., pharynx, rectum, urethra) and that testing include serologic screening for syphilis.[51] Any identified sexually transmitted infection should be promptly treated along with treatment of the partner.[52]

Prevention Strategies in Persons with Substance Use

Substance use disorders among people are a recognized cofactor for HIV transmission.[[53,54,55](#)] Substance use can impact sexual activity and can potentially lower rates of antiretroviral therapy adherence.[[56](#)] Alcohol use is the most prevalent risk factor for poor HIV medication adherence and lower rates of viral suppression.[[57,58](#)] Methamphetamine and other amphetamine-type stimulant use can alter antiretroviral medication adherence and increase sexual activities associated with increased risk of HIV transmission.[[59,60](#)] In addition, injection of methamphetamine and/or opioids can enhance the risk of HIV transmission through sharing of injection equipment. Screening for substance use disorders in people with HIV can play a significant role in prevention, especially if persons identified with a substance use disorder can receive counseling, treatment, and prevention services.[[61,62](#)]

Harm Reduction Approach

Harm reduction is based on a set of practical strategies and ideas aimed at reducing negative consequences associated with drug use. In addition, harm reduction programs do not insist on abstinence, and these programs typically utilize a spectrum of services, including providing safe injection counseling, HIV prevention education, and opiate substitution therapy. Harm reduction programs ideally provide a comprehensive set of services, including HIV counseling and testing (for individuals with HIV and their needle-sharing partners) and referral to substance use treatment programs. It is extremely important to remember that persons who inject drugs can also acquire and transmit HIV via sexual contact and should be counseled about sexual risk reduction strategies.[[63](#)]

Summary Points

- In the United States, integrated, evidence-based biomedical, behavioral, and structural interventions can substantially reduce transmission of HIV from persons with HIV to others.
- Antiretroviral therapy is recommended for all persons with HIV to prevent HIV transmission to others. Persons with HIV who consistently take antiretroviral therapy and maintain undetectable HIV RNA levels do not sexually transmit HIV to others, even with condomless sex.
- Persons unaware of their HIV status account for a disproportionate number of new HIV infections in the United States.
- For persons newly diagnosed with HIV, partner notification and contact HIV testing provides effective HIV case finding and the opportunity to decrease exposure to others.
- Consistent and correct condom use decreases HIV transmission by approximately 80% among serodifferent heterosexual couples and reduces the per-contact risk of HIV infection by 78% among men who have sex with men who practice receptive anal intercourse.
- Persons with undiagnosed acute (early) HIV have the highest relative risk of HIV transmission to others. Diagnosis, counseling, and treatment of persons with acute HIV can substantially reduce new HIV infections.
- Screening and treatment of sexually transmitted infections is an important component of overall HIV prevention services.
- Screening for substance use can help providers identify individuals with HIV who could benefit from counseling and support for a substance use disorder.

Citations

1. Centers for Disease Control and Prevention. Estimated HIV Incidence and Prevalence in the United States, 2018–2022. HIV Surveillance Supplemental Report. 2024;29(No. 1):1-131. Published May 2024 (revised February 7, 2025).
[\[CDC\]](#) -
2. Rodger AJ, Cambiano V, Bruun T, et al. Sexual Activity Without Condoms and Risk of HIV Transmission in Serodifferent Couples When the HIV-Positive Partner Is Using Suppressive Antiretroviral Therapy. JAMA. 2016;316:171-81.
[\[PubMed Abstract\]](#) -
3. Cohen MS, Chen YQ, McCauley M, et al. Prevention of HIV-1 infection with early antiretroviral therapy. N Engl J Med. 2011;365:493-505.
[\[PubMed Abstract\]](#) -
4. Cohen MS, Chen YQ, McCauley M, et al. Antiretroviral Therapy for the Prevention of HIV-1 Transmission. N Engl J Med. 2016;375:830-9.
[\[PubMed Abstract\]](#) -
5. Eisinger RW, Dieffenbach CW, Fauci AS. HIV Viral Load and Transmissibility of HIV Infection: Undetectable Equals Untransmittable. JAMA. 2019;321:451-2.
[\[PubMed Abstract\]](#) -
6. Panel on Antiretroviral Guidelines for Adults and Adolescents. Guidelines for the use of antiretroviral agents in adults and adolescents living with HIV. Department of Health and Human Services. Antiretroviral therapy to prevent sexual transmission of HIV (treatment as prevention). September 25, 2025
[\[HIV.gov\]](#) -
7. Quinn TC, Wawer MJ, Sewankambo N, et al. Viral load and heterosexual transmission of human immunodeficiency virus type 1. Rakai Project Study Group. N Engl J Med. 2000;342:921-9.
[\[PubMed Abstract\]](#) -
8. McNairy ML, El-Sadr WM. Antiretroviral therapy for the prevention of HIV transmission: what will it take? Clin Infect Dis. 2014;58:1003-11.
[\[PubMed Abstract\]](#) -
9. Attia S, Egger M, Müller M, Zwahlen M, Low N. Sexual transmission of HIV according to viral load and antiretroviral therapy: systematic review and meta-analysis. AIDS. 2009;23:1397-404.
[\[PubMed Abstract\]](#) -
10. Rodger AJ, Cambiano V, Bruun T, et al. Risk of HIV transmission through condomless sex in serodifferent gay couples with the HIV-positive partner taking suppressive antiretroviral therapy (PARTNER): final results of a multicentre, prospective, observational study. Lancet. 2019;393:2428-38.
[\[PubMed Abstract\]](#) -
11. Bavinton BR, Pinto AN, Phanuphak N, et al. Viral suppression and HIV transmission in serodiscordant male couples: an international, prospective, observational, cohort study. Lancet HIV. 2018;5:e438-e447.
[\[PubMed Abstract\]](#) -
12. Panel on Antiretroviral Guidelines for Adults and Adolescents. Guidelines for the use of antiretroviral

agents in adults and adolescents with HIV. Department of Health and Human Services. Initiation of antiretroviral therapy. September 25, 2025.

[\[HIV.gov\]](#) -

13. Hall HI, An Q, Tang T, et al. Prevalence of Diagnosed and Undiagnosed HIV Infection - United States, 2008-2012. *MMWR Morb Mortal Wkly Rep.* 2015;64:657-62.
[\[PubMed Abstract\]](#) -
14. Hall HI, Frazier EL, Rhodes P, et al. Differences in human immunodeficiency virus care and treatment among subpopulations in the United States. *JAMA Intern Med.* 2013;173:1337-44.
[\[Pub Med Abstract\]](#) -
15. Hall HI, Holtgrave DR, Maulsby C. HIV transmission rates from persons living with HIV who are aware and unaware of their infection. *AIDS.* 2012;26:893-6.
[\[PubMed Abstract\]](#) -
16. Marks G, Crepaz N, Janssen RS. Estimating sexual transmission of HIV from persons aware and unaware that they are infected with the virus in the USA. *AIDS.* 2006;20:1447-50.
[\[PubMed Abstract\]](#) -
17. Li Z, Purcell DW, Sansom SL, Hayes D, Hall HI. Vital Signs: HIV transmission along the continuum of care - United States, 2016. *MMWR Morb Mortal Wkly Rep.* 2019;68:267-72.
[\[PubMed Abstract\]](#) -
18. Fauci AS, Redfield RR, Sigounas G, Weahkee MD, Giroir BP. Ending the HIV Epidemic: A Plan for the United States. *JAMA.* 2019;321:844-845.
[\[PubMed Abstract\]](#) -
19. Sullivan PS, Carballo-Diéguez A, Coates T, et al. Successes and challenges of HIV prevention in men who have sex with men. *Lancet.* 2012;380:388-99.
[\[PubMed Abstract\]](#) -
20. Centers for Disease Control and Prevention, Health Resources and Services Administration, National Institutes of Health, American Academy of HIV Medicine, Association of Nurses in AIDS Care, International Association of Providers of AIDS Care, the National Minority AIDS Council, and Urban Coalition for HIV/AIDS Prevention Services. Recommendations for HIV Prevention with Adults and Adolescents with HIV in the United States, 2014. December 11, 2014 (amended December 30, 2016)
[\[CDC\]](#) -
21. Crepaz N, Lyles CM, Wolitski RJ, et al. Do prevention interventions reduce HIV risk behaviours among people living with HIV? A meta-analytic review of controlled trials. *AIDS.* 2006;20:143-57.
[\[PubMed Abstract\]](#) -
22. Golden MR, Dombrowski JC, Wood RW, Fleming M, Harrington RD. A controlled study of the effectiveness of public health HIV partner notification services. *AIDS.* 2009;23:133-5.
[\[PubMed Abstract\]](#) -
23. Hogben M, McNally T, McPheeters M, Hutchinson AB. The effectiveness of HIV partner counseling and referral services in increasing identification of HIV-positive individuals a systematic review. *Am J Prev Med.* 2007;33:S89-100.
[\[PubMed Abstract\]](#) -
24. Golden MR, Hogben M, Potterat JJ, Handsfield HH. HIV partner notification in the United States: a national survey of program coverage and outcomes. *Sex Transm Dis.* 2004;31:709-12.

[\[PubMed Abstract\]](#) -

25. Katz DA, Hogben M, Dooley SW Jr, Golden MR. Increasing public health partner services for human immunodeficiency virus: results of a second national survey. *Sex Transm Dis.* 2010;37:469-75.
[\[PubMed Abstract\]](#) -
26. Weller S, Davis K. Condom effectiveness in reducing heterosexual HIV transmission. *Cochrane Database Syst Rev.* 2002;:CD003255.
[\[PubMed Abstract\]](#) -
27. Giannou FK, Tsiara CG, Nikolopoulos GK, et al. Condom effectiveness in reducing heterosexual HIV transmission: a systematic review and meta-analysis of studies on HIV serodiscordant couples. *Expert Rev Pharmacoecon Outcomes Res.* 2016;16:489-99.
[\[PubMed Abstract\]](#) -
28. Hughes JP, Baeten JM, Lingappa JR, et al. Determinants of per-coital-act HIV-1 infectivity among African HIV-1-serodiscordant couples. *J Infect Dis.* 2012;205:358-65.
[\[PubMed Abstract\]](#) -
29. Smith DK, Herbst JH, Zhang X, Rose CE. Condom effectiveness for HIV prevention by consistency of use among men who have sex with men in the United States. *J Acquir Immune Defic Syndr.* 2015;68:337-44.
[\[PubMed Abstract\]](#) -
30. Cassels S, Katz DA. Seroadaptation among men who have sex with men: emerging research themes. *Curr HIV/AIDS Rep.* 2013;10:305-13.
[\[PubMed Abstract\]](#) -
31. Kennedy CE, Bernard LJ, Muessig KE, et al. *J Sex Transm Dis.* 2013;2013:583627.
[\[PubMed Abstract\]](#) -
32. Truong HM, Kellogg T, Klausner JD, et al. Increases in sexually transmitted infections and sexual risk behaviour without a concurrent increase in HIV incidence among men who have sex with men in San Francisco: a suggestion of HIV serosorting? *Sex Transm Infect.* 2006;82:461-6.
[\[PubMed Abstract\]](#) -
33. van den Boom W, Konings R, Davidovich U, Sandfort T, Prins M, Stolte IG. Is serosorting effective in reducing the risk of HIV infection among men who have sex with men with casual sex partners? *J Acquir Immune Defic Syndr.* 2014;65:375-9.
[\[PubMed Abstract\]](#) -
34. Khosropour CM, Dombrowski JC, Swanson F, et al. Trends in Serosorting and the Association With HIV/STI Risk Over Time Among Men Who Have Sex With Men. *J Acquir Immune Defic Syndr.* 2016;72:189-97.
[\[PubMed Abstract\]](#) -
35. Cohen MS, Muessig KE, Smith MK, Powers KA, Kashuba AD. Antiviral agents and HIV prevention: controversies, conflicts, and consensus. *AIDS.* 2012;26:1585-98.
[\[PubMed Abstract\]](#) -
36. Pilcher CD, Joaki G, Hoffman IF, et al. Amplified transmission of HIV-1: comparison of HIV-1 concentrations in semen and blood during acute and chronic infection. *AIDS.* 2007;21:1723-30.
[\[PubMed Abstract\]](#) -

37. Pilcher CD, Tien HC, Eron JJ Jr, et al. Brief but efficient: acute HIV infection and the sexual transmission of HIV. *J Infect Dis.* 2004;189:1785-92.
[\[PubMed Abstract\]](#) -
38. Hollingsworth TD, Anderson RM, Fraser C. HIV-1 transmission, by stage of infection. *J Infect Dis.* 2008;198:687-93.
[\[PubMed Abstract\]](#) -
39. Wawer MJ, Gray RH, Sewankambo NK, et al. Rates of HIV-1 transmission per coital act, by stage of HIV-1 infection, in Rakai, Uganda. *J Infect Dis.* 2005;191:1403-9.
[\[PubMed Abstract\]](#) -
40. Brenner BG, Roger M, Routy JP, et al. High rates of forward transmission events after acute/early HIV-1 infection. *J Infect Dis.* 2007;195:951-9.
[\[PubMed Abstract\]](#) -
41. Powers KA, Ghani AC, Miller WC, et al. The role of acute and early HIV infection in the spread of HIV and implications for transmission prevention strategies in Lilongwe, Malawi: a modelling study. *Lancet.* 2011;378:256-68.
[\[PubMed Abstract\]](#) -
42. Centers for Disease Control and Prevention (CDC). National HIV Testing Day and new testing recommendations. *MMWR Morb Mortal Wkly Rep.* 2014;63:537.
[\[PubMed Abstract\]](#) -
43. Centers for Disease Control and Prevention and Association of Public Health Laboratories. Laboratory testing for the diagnosis of HIV infection: updated recommendations. June 27, 2014.
[\[CDC\]](#) -
44. Hamlyn E, Jones V, Porter K, Fidler S. Antiretroviral treatment of primary HIV infection to reduce onward transmission. *Curr Opin HIV AIDS.* 2010;5:283-90.
[\[PubMed Abstract\]](#) -
45. Panel on Antiretroviral Guidelines for Adults and Adolescents. Guidelines for the Use of Antiretroviral Agents in Adults and Adolescents with HIV. Department of Health and Human Services. Considerations for Antiretroviral Therapy Use in Special Populations. Early (Acute and Recent) HIV Infection. September 12, 2024.
[\[HIV.gov\]](#) -
46. Marrazzo JM, del Rio C, Holtgrave DR, et al. HIV prevention in clinical care settings: 2014 recommendations of the International Antiviral Society-USA Panel. *JAMA.* 2014;312:390-409.
[\[PubMed Abstract\]](#) -
47. Mayer KH. Introduction: Linkage, engagement, and retention in HIV care: essential for optimal individual- and community-level outcomes in the era of highly active antiretroviral therapy. *Clin Infect Dis.* 2011;52 Suppl 2:S205-7.
[\[PubMed Abstract\]](#) -
48. Garg S, Taylor LE, Grasso C, Mayer KH. Prevalent and incident hepatitis C virus infection among HIV-infected men who have sex with men engaged in primary care in a Boston community health center. *Clin Infect Dis.* 2013;56:1480-7.
[\[PubMed Abstract\]](#) -
49. Spindler HH, Scheer S, Chen SY, et al. Viagra, methamphetamine, and HIV risk: results from a

probability sample of MSM, San Francisco. Sex Transm Dis. 2007;34:586-91.

[\[PubMed Abstract\]](#) -

50. Celum C, Wald A, Lingappa JR, et al. Acyclovir and transmission of HIV-1 from persons infected with HIV-1 and HSV-2. N Engl J Med. 2010;362:427-39.

[\[PubMed Abstract\]](#) -

51. Workowski KA, Bachmann LH, Chan PA, et al. Sexually transmitted infections treatment guidelines, 2021. Screening recommendations referenced in treatment guidelines and original recommendation sources. MMWR Recomm Rep. 2021;70(No. RR-4):1-187.

[\[2021 STD Treatment Guidelines\]](#) -

52. Workowski KA, Bachmann LH, Chan PA, et al. Sexually transmitted infections treatment guidelines, 2021. MMWR Recomm Rep. 2021;70(No. RR-4):1-187.

[\[2021 STI Treatment Guidelines\]](#) -

53. Morin SF, Myers JJ, Shade SB, Koester K, Maiorana A, Rose CD. Predicting HIV transmission risk among HIV-infected patients seen in clinical settings. AIDS Behav. 2007;11:S6-16.

[\[PubMed Abstract\]](#) -

54. Mimiaga MJ, Reisner SL, Grasso C, et al. Substance use among HIV-infected patients engaged in primary care in the United States: findings from the Centers for AIDS Research Network of Integrated Clinical Systems cohort. Am J Public Health. 2013;103:1457-67.

[\[PubMed Abstract\]](#) -

55. Samet JH, Cheng DM, Libman H, Nunes DP, Alperen JK, Saitz R. Alcohol consumption and HIV disease progression. J Acquir Immune Defic Syndr. 2007;46:194-9.

[\[PubMed Abstract\]](#) -

56. Clark T, Marquez C, Hare CB, John MD, Klausner JD. Methamphetamine use, transmission risk behavior and internet use among HIV-infected patients in medical care, San Francisco, 2008. AIDS Behav. 2012;16:396-403.

[\[PubMed Abstract\]](#) -

57. Kalichman SC, Grebler T, Amaral CM, et al. Intentional non-adherence to medications among HIV positive alcohol drinkers: prospective study of interactive toxicity beliefs. J Gen Intern Med. 2013;28:399-405.

[\[PubMed Abstract\]](#) -

58. Kalichman SC, Grebler T, Amaral CM, et al. Viral suppression and antiretroviral medication adherence among alcohol using HIV-positive adults. Int J Behav Med. 2014;21:811-20.

[\[PubMed Abstract\]](#) -

59. Forrest DW, Metsch LR, LaLota M, Cardenas G, Beck DW, Jeanty Y. Crystal methamphetamine use and sexual risk behaviors among HIV-positive and HIV-negative men who have sex with men in South Florida. J Urban Health. 2010;87:480-5.

[\[PubMed Abstract\]](#) -

60. Semple SJ, Zians J, Grant I, Patterson TL. Sexual risk behavior of HIV-positive methamphetamine-using men who have sex with men: the role of partner serostatus and partner type. Arch Sex Behav. 2006;35:461-71.

[\[PubMed Abstract\]](#) -

61. Aspinall EJ, Nambiar D, Goldberg DJ, et al. Are needle and syringe programmes associated with a

reduction in HIV transmission among people who inject drugs: a systematic review and meta-analysis. *Int J Epidemiol.* 2014;43:235-48.

[\[PubMed Abstract\]](#) -

62. Centers for Disease Control and Prevention (CDC). Syringe exchange programs --- United States, 2008. *MMWR Morb Mortal Wkly Rep.* 2010;59:1488-91.

[\[PubMed Abstract\]](#) -

63. Kral AH, Bluthenthal RN, Lorvick J, Gee L, Bacchetti P, Edlin BR. Sexual transmission of HIV-1 among injection drug users in San Francisco, USA: risk-factor analysis. *Lancet.* 2001;357:1397-401.

[\[PubMed Abstract\]](#) -

References

- Boily MC, Baggaley RF, Wang L, et al. Heterosexual risk of HIV-1 infection per sexual act: systematic review and meta-analysis of observational studies. *Lancet Infect Dis.* 2009;9:118-29.
[\[PubMed Abstract\]](#) -
- Branson BM, Handsfield HH, Lampe MA, et al. Revised recommendations for HIV testing of adults, adolescents, and pregnant women in health-care settings. *MMWR Recomm Rep.* 2006;55:1-17.
[\[PubMed Abstract\]](#) -
- Castel AD, Befus M, Willis S, et al. Use of the community viral load as a population-based biomarker of HIV burden. *AIDS.* 2012;26:345-53.
[\[PubMed Abstract\]](#) -
- Centers for Disease Control and Prevention. Monitoring selected national HIV prevention and care objectives by using HIV surveillance data--United States and 6 U.S. dependent areas, 2014. *HIV Surveillance Supplemental Report.* 2016;21(No. 4):1-87. Published July 2016.
[\[CDC\]](#) -
- Crosby RA, Mena L, Geter A. Favourable attitudes towards serosorting are associated with overall less frequent condom use among young Black men having sex with men. *Sex Health.* 2016;13:91-2.
[\[PubMed Abstract\]](#) -
- Das M, Chu PL, Santos GM, et al. Decreases in community viral load are accompanied by reductions in new HIV infections in San Francisco. *PLoS One.* 2010;5:e11068.
[\[PubMed Abstract\]](#) -
- Fisher M, Pao D, Brown AE, et al. Determinants of HIV-1 transmission in men who have sex with men: a combined clinical, epidemiological and phylogenetic approach. *AIDS.* 2010;24:1739-47.
[\[PubMed Abstract\]](#) -
- Galvin SR, Cohen MS. The role of sexually transmitted diseases in HIV transmission. *Nat Rev Microbiol.* 2004;2:33-42.
[\[PubMed Abstract\]](#) -
- Holt M, Lea T, Schmidt HM, et al. Increasing Belief in the Effectiveness of HIV Treatment as Prevention: Results of Repeated, National Surveys of Australian Gay and Bisexual Men, 2013-15. *AIDS Behav.* 2016;20:1564-71.
[\[PubMed Abstract\]](#) -
- Hoover KW, Butler M, Workowski K, et al. STD screening of HIV-infected MSM in HIV clinics. *Sex*

Transm Dis. 2010;37:771-6.

[\[PubMed Abstract\]](#) -

- Koff A, Goldberg C, Ogbuagu O. Condomless sex and HIV transmission among serodifferent couples: current evidence and recommendations. *Ann Med.* 2017;49:534-544.
[\[PubMed Abstract\]](#) -
- Marks G, Crepaz N, Senterfitt JW, Janssen RS. Meta-analysis of high-risk sexual behavior in persons aware and unaware they are infected with HIV in the United States: implications for HIV prevention programs. *J Acquir Immune Defic Syndr.* 2005;39:446-53.
[\[PubMed Abstract\]](#) -
- Miller WC, Powers KA, Smith MK, Cohen MS. Community viral load as a measure for assessment of HIV treatment as prevention. *Lancet Infect Dis.* 2013;13:459-64.
[\[PubMed Abstract\]](#) -
- Montaner JS, Lima VD, Barrios R, et al. Association of highly active antiretroviral therapy coverage, population viral load, and yearly new HIV diagnoses in British Columbia, Canada: a population-based study. *Lancet.* 2010;376:532-9.
[\[PubMed Abstract\]](#) -
- Patton ME, Kidd S, Llata E, et al. Extragenital gonorrhea and chlamydia testing and infection among men who have sex with men--STD Surveillance Network, United States, 2010-2012. *Clin Infect Dis.* 2014;58:1564-70.
[\[PubMed Abstract\]](#) -
- Smith MK, Rutstein SE, Powers KA, et al. The detection and management of early HIV infection: a clinical and public health emergency. *J Acquir Immune Defic Syndr.* 2013;63 Suppl 2:S187-99.
[\[PubMed Abstract\]](#) -
- Van Howe RS. Genital ulcerative disease and sexually transmitted urethritis and circumcision: a meta-analysis. *Int J STD AIDS.* 2007;18:799-809.
[\[PubMed Abstract\]](#) -
- Wilson DP, Regan DG, Heymer KJ, Jin F, Prestage GP, Grulich AE. Serosorting may increase the risk of HIV acquisition among men who have sex with men. *Sex Transm Dis.* 2010;37:13-7.
[\[PubMed Abstract\]](#) -

Figures

Figure 1 Estimated HIV Incidence in United States, 2018-2022

Source: Centers for Disease Control and Prevention. Estimated HIV Incidence and Prevalence in the United States, 2018–2022. HIV Surveillance Supplemental Report. 2024;29(No. 1):1-131. Published May 2024 (revised February 7, 2025).




Figure 2 Adjusted Rate Ratio of Heterosexual Transmission of HIV-1 According to Serum HIV-1 RNA Level of the HIV-1 Positive Partner

This graphic illustrates the correlation of risk of HIV transmission and serum HIV-1 levels in the person infected with HIV. No HIV transmissions occurred from persons with HIV who had serum HIV RNA-1 levels less than 1,500 copies/mL.

Source: Quinn TC, Wawer MJ, Sewankambo N, et al. Viral load and heterosexual transmission of human immunodeficiency virus type 1. Rakai Project Study Group. N Engl J Med. 2000;342:921-9.




Figure 3 (Image Series) - HPTN 052 and Antiretroviral Therapy for the Prevention of HIV-1
(Image Series) - Figure 3 (Image Series) - HPTN 052 and Antiretroviral Therapy for the Prevention of HIV-1

Image 3A: HPTN 052 Patient Population

The HPTN 052 trial enrolled 1,763 HIV serodifferent couples and 97% identified as heterosexual couples.

Source: Cohen MS, Chen YQ, McCauley M, et al. Prevention of HIV-1 infection with early antiretroviral therapy. N Engl J Med. 2011;365:493-505.

1,763 HIV Serodifferent Couples (97% heterosexual)




Figure 3 (Image Series) - HPTN 052 and Antiretroviral Therapy for the Prevention of HIV-1
Image 3B: HPTN 052 Study Design

Adults with HIV in the early therapy arm received combination antiretroviral therapy and those in the deferred therapy arm started antiretroviral therapy when their CD4 decreased to less than 250 cells/mm³ or they had an AIDS-related event.

Source: Cohen MS, Chen YQ, McCauley M, et al. Prevention of HIV-1 infection with early antiretroviral therapy. N Engl J Med. 2011;365:493-505.




Figure 3 (Image Series) - HPTN 052 and Antiretroviral Therapy for the Prevention of HIV-1
Image 3C: HPTN 052 Results

This graphic shows linked transmissions. Couples in the early therapy arm had a 96% reduction in new HIV transmission events.

Source: Cohen MS, Chen YQ, McCauley M, et al. Prevention of HIV-1 infection with early antiretroviral therapy. N Engl J Med. 2011;365:493-505.




Figure 4 Transmission of HIV and Awareness of HIV Status

This graph shows the estimated percentage of HIV transmissions based on awareness of HIV diagnosis. These estimates were based on the 2016 Center for Disease Control and Prevention (CDC) Progression and Transmission of HIV (PATH 2.0) model.

Source: Li Z, Purcell DW, Sansom SL, Hayes D, Hall HI. Vital Signs: HIV transmission along the continuum of care - United States, 2016. MMWR Morb Mortal Wkly Rep. 2019;68:267-72.




Figure 5 Risk of Heterosexual Male HIV Acquisition Based on Circumcision Status of Males Living in Africa

Abbreviation: RR = relative risk

These three studies conducted in Africa addressed the risk of heterosexual HIV acquisition in men based on their circumcision status. As shown, the risk of HIV acquisition was significantly lower in men who were circumcised when compared with those who were uncircumcised.

Source: (1) Auvert B, Taljaard D, Lagarde E, Sobngwi-Tambekou J, Sitta R, Puren A. Randomized, controlled intervention trial of male circumcision for reduction of HIV infection risk: the ANRS 1265 Trial. *PLoS Med.* 2005;2:e298. / (2) Bailey RC, Moses S, Parker CB, et al. Male circumcision for HIV prevention in young men in Kisumu, Kenya: a randomised controlled trial. *Lancet.* 2007;369:643-56. / (3) Gray RH, Kigozi G, Serwadda D, et al. Male circumcision for HIV prevention in men in Rakai, Uganda: a randomised trial. *Lancet.* 2007;369:657-66.




Figure 6 Relative Risk of HIV Transmission Along the HIV Care Continuum

This graph shows the estimated transmission rate based on transmissions per 100,000 person-years. These estimates were based on the 2016 Center for Disease Control and Prevention (CDC) Progression and Transmission of HIV (PATH 2.0) model.

Source: Li Z, Purcell DW, Sansom SL, Hayes D, Hall HI. Vital Signs: HIV transmission along the continuum of care - United States, 2016. MMWR Morb Mortal Wkly Rep. 2019;68:267-72.




Figure 7 Risk of Heterosexual Female HIV Acquisition Based on Circumcision Status of Male Sex Partner

In this study, investigators in Rakai, Uganda enrolled 922 uncircumcised males with HIV infection who were randomized to undergo immediate circumcision (intervention group) or have circumcision delayed for 24 months (control group). The trial was stopped early because of futility and there was no reduction in risk of female HIV acquisition from their male partners who had been circumcised.

Source: Wawer MJ, Makumbi F, Kigozi G, et al. Circumcision in HIV-infected men and its effect on HIV transmission to female partners in Rakai, Uganda: a randomised controlled trial. *Lancet*. 2009;374:229-37.




Table 1. Antiretroviral Therapy to Prevent Sexual Transmission of HIV (Treatment as Prevention)

<p>Guidelines for the Use of Antiretroviral Agents in Adults and Adolescents with HIV</p> <p>Use of Antiretroviral Therapy to Prevent Sexual Transmission of HIV</p> <ul style="list-style-type: none"> • All people with HIV should be informed that maintaining a plasma HIV RNA (viral load) of <200 copies/mL, including any measurable value below this threshold value, with antiretroviral therapy (ART) prevents sexual transmission of HIV. This concept may be recognized as Undetectable = Untransmittable (AII). • People with HIV who are starting ART should use another form of prevention with sexual partners (e.g. condoms, pre-exposure prophylaxis [PrEP] for the sexual partner who is HIV-negative) for at least the first 6 months of ART and until a viral load of <200 copies/mL has been documented (AII). Many experts would recommend confirming sustained suppression before assuming that there is no further risk of sexual HIV transmission (AIII). • When the viral load is ≥ 200 copies/mL, at least one additional HIV prevention method (e.g., condoms, sexual abstinence, or HIV PrEP for the HIV-negative partner) is recommended to prevent sexual HIV transmission until resuppression to <200 copies/mL is confirmed (AIII). • People with HIV who intend to rely upon ART for prevention need to maintain high levels of ART adherence (AIII). They should be informed that transmission is possible during periods of poor adherence or treatment interruption (AIII). • At each visit for HIV care, clinicians should assess adherence to ART and counsel people with HIV on the importance of ART for their own health as well as its role in preventing sexual HIV transmission (AIII). • Providers should inform patients that maintaining a viral load of <200 copies/mL does not prevent acquisition or transmission of other sexually transmitted infections (STIs) (AII). • Providers should also routinely screen all people with HIV who are sexually active for STIs, both for their own health and to prevent transmission of STIs to others (AIII). <p>Rating of Recommendations: A = Strong; B = Moderate; C = Optional</p> <p>Rating of Evidence: I = Data from randomized controlled trials; II = Data from well-designed nonrandomized trials or observational cohort studies with long-term clinical outcomes; III = Expert opinion</p>

Source:

- Panel on Antiretroviral Guidelines for Adults and Adolescents. Guidelines for the use of antiretroviral agents in adults and adolescents living with HIV. Department of Health and Human Services. Antiretroviral therapy to prevent sexual transmission of HIV (treatment as prevention). September 25, 2025 [[HIV.gov](https://www.hiv.gov)]

